

8th Homework Set — Solutions

Chapter 6

Problem 6.11 Let A be the number of people buying an ordinary set, B the number of people buying a plasma set, and C the number of people who are just browsing. Then $P\{A = 2, B = 1, C = 2\} = \frac{5!}{2!1!2!} 0.45^2 \cdot 0.15 \cdot 0.4^2 = 0.1458$.

Problem 6.13 Let X be uniform on $(-15, 15)$, and let Y be uniform on $(-30, 30)$. Nobody waits longer than five minutes if $|Y - X| < 5$.

$$\begin{aligned} P\{|Y - X| < 5\} &= P\{-5 < Y - X < 5\} \\ &= P\{X - 5 < Y < X + 5\} \\ &= \int_{-15}^{15} \int_{x-5}^{x+5} \frac{1}{30 \cdot 60} dy dx \\ &= \frac{30 \cdot 10}{30 \cdot 60} = \frac{1}{6}. \end{aligned}$$

The probability that the man arrives first is $P\{X < Y\} = \frac{1}{2}$ by symmetry.

Problem 6.14 Let X, Y be uniform random variables on $(0, L)$. Let $Z = |Y - X|$. We want to find $E[Z]$. First, find $F_Z(a)$, for $a \geq 0$. We have $F_Z(a) = P\{Z \leq a\} = P\{|Y - X| \leq a\} = P\{-a \leq Y - X \leq a\} = \frac{2aL - a^2}{L^2}$. using geometric considerations. Hence, $f_Z(x) = \frac{2L - 2x}{L^2}$ if $0 \leq a \leq L$. Hence,

$$\begin{aligned} E[Z] &= \int_0^L x \cdot \frac{2L - 2x}{L^2} dx \\ &= \frac{2}{L^2} \left(\frac{Lx^2}{2} - \frac{x^3}{3} \right) \Big|_0^L \\ &= \frac{L}{3}. \end{aligned}$$

Problem 6.18 Let X be uniform on $(0, \frac{L}{2})$ and let Y be uniform on $(\frac{L}{2}, L)$. We want

to find $P\{Y - X > \frac{L}{3}\}$.

$$\begin{aligned} P\left\{Y - X > \frac{L}{3}\right\} &= P\left\{Y < \frac{L}{2} + \frac{L}{3}, X < Y - \frac{L}{3}\right\} + P\left\{Y > \frac{L}{2} + \frac{L}{3}\right\} \\ &= \int_{\frac{L}{2}}^{\frac{5L}{6}} \int_0^{y-\frac{L}{3}} \frac{4}{L^2} dx dy + \int_{\frac{5L}{6}}^{\frac{L}{2}} \frac{2}{L} dy \\ &= \frac{4}{9} + \frac{1}{3} = \frac{7}{9}. \end{aligned}$$

Problem 6.20 If the joint density function of X and Y is

$$f(x, y) = \begin{cases} xe^{-(x+y)} & x > 0, y > 0 \\ 0 & \text{otherwise,} \end{cases}$$

then $f(x, y) = f_X(x)f_Y(y)$, where $f_X(x) = xe^{-x}$ for $x > 0$, and $f_Y(y) = e^{-y}$ for $y > 0$ (0 otherwise), so that X and Y are independent.

If

$$f(x, y) = \begin{cases} 2 & 0 < x < y, 0 < y < 1 \\ 0 & \text{otherwise,} \end{cases}$$

then X and Y are not independent because the nonzero values of f are located in a triangular domain.

- Problem 6.21 (a) Check: $\int_0^1 \int_0^{1-y} 24xy dx dy = \int_0^1 12(1-y)^2 y dy = 12 \int_0^1 y - 2y^2 + y^3 dy = 6y^2 - 8y^3 + 3y^4 \Big|_0^1 = 6 - 8 + 3 = 1$.
- (b) First, find $f_X(x) = \int_0^{1-x} 24xy dy = 12x(1-x)^2$. Now, $E[X] = \int_0^1 12x^2(1-x)^2 dx = 4x^2 - 6x^3 + \frac{12}{5}x^5 \Big|_0^1 = 4 - 6 + \frac{12}{5} = \frac{2}{5}$.
- (c) $E[Y] = E[X] = \frac{2}{5}$ by symmetry.

Problem 6.22 Let X and Y be jointly continuous with density function

$$f(x, y) = \begin{cases} x + y & 0 < x < 1, 0 < y < 1 \\ 0 & \text{otherwise.} \end{cases}$$

- (a) X and Y are not independent, since $f(x, y)$ is clearly not a product of functions of x and y .
- (b) $f_X(x) = \int_0^1 x + y dy = x + \frac{y^2}{2} \Big|_0^1 = x + \frac{1}{2}$.

$$(c) P\{X + Y < 1\} = \int_0^1 \int_0^{1-y} x + y dx dy = \int_0^1 \frac{(1-y)^2}{2} + y(1-y) dy = \frac{1}{2} \int_0^1 1 - y^2 dy = \frac{1}{2} \left(1 - \frac{1}{3}\right) = \frac{1}{3}.$$

Problem 6.23 Let X and Y be jointly distributed with density function

$$f(x, y) = \begin{cases} 12xy(1-x) & 0 < x < 1, 0 < y < 1 \\ 0 & \text{otherwise.} \end{cases}$$

First, compute $f_X(x) = \int_0^1 12xy(1-x) dy = 6x(1-x)$ and $f_Y(y) = \int_0^1 12xy(1-x) dx = 2y$.

- (a) Clearly, $f(x, y) = f_X(x)f_Y(y)$, so that X and Y are independent.
- (b) $E[X] = \int_0^1 6x^2(1-x) dx = 2x^3 - \frac{3}{2}x^4 \Big|_0^1 = \frac{1}{2}$.
- (c) $E[Y] = \int_0^1 2y^2 dy = \frac{2}{3}y^3 \Big|_0^1 = \frac{2}{3}$.
- (d) First, find $E[X^2] = \int_0^1 6x^3(1-x) dx = \frac{3}{2}x^4 - \frac{6}{5}x^5 \Big|_0^1 = \frac{3}{10}$. Now, $\text{Var}(X) = E[X^2] - E[X]^2 = \frac{3}{10} - \frac{1}{4} = \frac{1}{20}$.
- (e) First, find $E[Y^2] = \int_0^1 2y^3 dy = \frac{1}{2}y^4 \Big|_0^1 = \frac{1}{2}$. Now, $\text{Var}(Y) = \frac{1}{2} - \frac{4}{9} = \frac{1}{18}$.

Problem 6.27 Let X_1, X_2 be exponential random variables with parameter λ_1, λ_2 . Let $Z = \frac{X_1}{X_2}$. Note that $F_Z(a) = 0$ if $a \leq 0$. Compute $F_Z(a)$ for $a > 0$:

$$\begin{aligned} F_Z(a) &= P\{Z \leq a\} = P\{X_1 \leq aX_2\} \\ &= \lambda_1 \lambda_2 \int_0^\infty \int_0^{ay} e^{-\lambda_1 x - \lambda_2 y} dx dy \\ &= \frac{\lambda_1 a}{\lambda_1 a + \lambda_2}, \end{aligned}$$

so that

$$f_Z(a) = \frac{d}{da} F(a) = \frac{\lambda_1}{\lambda_1 a + \lambda_2} - \frac{\lambda_1^2 a}{(a\lambda_1 + \lambda_2)^2}.$$

Finally, we have

$$P\{X_1 < X_2\} = P\{Z < 1\} = F_Z(1) = \frac{\lambda_1}{\lambda_1 + \lambda_2}.$$

Problem 6.29 Let X_1, X_2 be independent normal random variables with $\mu = 2200$ and $\sigma^2 = 230^2$, representing the gross sales over this week and next week, respectively. Then $X = X_1 + X_2$ is normal with mean 4400 and variance $2 \cdot 230^2 = 105800$.

$$(a) P\{X > 5000\} = P\left\{\frac{X-4400}{\sqrt{105800}} > \frac{600}{\sqrt{105800}}\right\} = 1 - \Phi(1.84) = 1 - 0.9671 = 0.0329.$$

$$(b) \text{ Let } p = P\{X_1 > 2000\} = P\left\{\frac{X_1-2200}{230} > \frac{-200}{230}\right\} = 1 - \Phi\left(-\frac{20}{23}\right) = \Phi(0.87) = 0.8078.$$

Let N be the number of weeks (out of three) in which the sales exceed \$2000. Then N is binomial with parameters $(p, 3)$, so that $P\{N \geq 2\} = p^3 + 3p^2(1-p) = 0.9034$.

Problem 6.31 Let X be the number of women who never eat breakfast, and let Y be the number of men who never eat breakfast. Let $Z = X + Y$. By DeMoivre-Laplace, X is approximated by a normal random variable with mean $200 \cdot 0.236 = 47.2$ and variance $47.2 \cdot 0.764 = 36.061$, and Y is normal with mean $200 \cdot 0.252 = 50.4$ and variance $50.4 \cdot 0.748 = 37.699$.

Let $Z_1 = X + Y$ and $Z_2 = X - Y$. Then Z_1 is normal with mean 97.6 and variance $36.061 + 37.699 = 73.76$, and Z_2 is normal with mean -3.2 and variance 73.76.

$$(a) P\{Z_1 \geq 110\} = P\{Z_1 > 109.5\} = P\left\{\frac{Z_1-97.6}{\sqrt{73.76}} > \frac{11.9}{\sqrt{73.76}}\right\} = 1 - \Phi(1.39) = 1 - 0.9177 = 0.0823.$$

$$(b) P\{X \geq Y\} = P\{X - Y \geq 0\} = P\{Z_2 \geq 0\} = P\{Z_2 > -0.5\} = P\left\{\frac{Z_2+3.2}{\sqrt{73.76}} > \frac{2.7}{\sqrt{73.76}}\right\} = 1 - \Phi(0.31) = 0.3783.$$

Problem 6.34 Let X_1 be the number of accidents in the next month, X_2 the number of accidents in the month after that, and X_3 the number of accidents in the third month. It makes sense to think of X_1, X_2 , and X_3 as independent Poisson random variables with parameter $\lambda = 2.2$.

Let $X = X_1$, $Y = X_1 + X_2$, and $Z = X_1 + X_2 + X_3$. Then X, Y , and Z are Poisson with parameter 2.2, 4.4, and 6.6, respectively.

$$(a) P\{X > 2\} = 1 - e^{-2.2} \left(1 + 2.2 + \frac{2.2^2}{2}\right) = 0.3773.$$

$$(b) P\{Y > 4\} = 1 - e^{-4.4} \left(1 + 4.4 + \frac{4.4^2}{2} + \frac{4.4^3}{3!} + \frac{4.4^4}{4!}\right) = 0.4488.$$

(c) $P\{Z > 5\} = 1 - e^{-6.6} \left(1 + 6.6 + \frac{6.6^2}{2} + \frac{6.6^3}{3!} + \frac{6.6^4}{4!} + \frac{6.6^5}{5!} \right) = 0.6453.$

Problem 6.38 (a) $P\{X = i, Y = j\} = \frac{1}{5^i}$ for $i = 1, \dots, 5$ and $j = 1, \dots, i$, 0 otherwise.

$P\{X = i, Y = j\}$	Y=1	Y=2	Y=3	Y=4	Y=5	$P\{X = i\}$
X=1	$\frac{1}{5}$	0	0	0	0	$\frac{1}{5}$
X=2	$\frac{1}{10}$	$\frac{1}{10}$	0	0	0	$\frac{1}{5}$
X=3	$\frac{1}{15}$	$\frac{1}{15}$	$\frac{1}{15}$	0	0	$\frac{1}{5}$
X=4	$\frac{1}{20}$	$\frac{1}{20}$	$\frac{1}{20}$	$\frac{1}{20}$	0	$\frac{1}{5}$
X=5	$\frac{1}{25}$	$\frac{1}{25}$	$\frac{1}{25}$	$\frac{1}{25}$	$\frac{1}{25}$	$\frac{1}{5}$
$P\{Y = j\}$	$\frac{137}{300}$	$\frac{77}{300}$	$\frac{47}{300}$	$\frac{9}{100}$	$\frac{1}{25}$	1

(b) $P\{X = i|Y = j\} = \frac{\frac{1}{5^i}}{\sum_{k=i}^5 \frac{1}{5^k}}$

$P\{X = i Y = j\}$	Y=1	Y=2	Y=3	Y=4	Y=5
X = 1	$\frac{60}{137}$	0	0	0	0
X = 2	$\frac{30}{137}$	$\frac{30}{77}$	0	0	0
X = 3	$\frac{20}{137}$	$\frac{20}{77}$	$\frac{20}{47}$	0	0
X = 4	$\frac{15}{137}$	$\frac{15}{77}$	$\frac{15}{47}$	$\frac{5}{9}$	0
X = 5	$\frac{12}{137}$	$\frac{12}{77}$	$\frac{12}{47}$	$\frac{4}{9}$	1

(c) No.

Problem 6.40

$p(i, j)$	$j = 1$	$j = 2$	
$i = 1$	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{3}{8}$
$i = 2$	$\frac{1}{8}$	$\frac{1}{2}$	$\frac{5}{8}$
	$\frac{1}{4}$	$\frac{3}{4}$	1

(a)

$P\{X = i Y = j\}$	$j = 1$	$j = 2$
$i = 1$	$\frac{1}{2}$	$\frac{1}{3}$
$i = 2$	$\frac{1}{2}$	$\frac{2}{3}$

(b) No.

(c) $P\{XY \leq 3\} = 1 - p(2, 2) = \frac{1}{2}$
 $P\{X + Y > 2\} = 1 - p(1, 1) = \frac{7}{8}$
 $P\{\frac{X}{Y} > 1\} = p(2, 1) = \frac{1}{8}$

Problem 6.41 Let X and Y be jointly continuous with density function $f(x, y) = xe^{-x(y+1)}$ for $x > 0, y > 0$. Note that $f_X(x) = \int_0^\infty f(x, y)dy = e^{-x}$ for $x > 0$, and $f_Y(y) = \int_0^\infty f(x, y)dx = \frac{1}{(y+1)^2}$ for $y > 0$.

(a) $f_{X|Y}(x|y) = (y+1)^2xe^{-x(y+1)}$ for $x > 0, y > 0$, 0 otherwise, and $f_{Y|X}(y|x) = xe^{-xy}$ for $x > 0, y > 0$.

(b) Let $Z = XY$. Then for $a > 0$,

$$F_Z(a) = P\{XY < a\} = \int_0^\infty \int_0^{\frac{a}{x}} xe^{-x(y+1)} dy dx = 1 - e^{-a}.$$

Hence, $f_Z(a) = \frac{d}{da}F_Z(a) = e^{-a}$ for $a > 0$, 0 otherwise.

Problem 6.42 Let X and Y be jointly continuous with density function

$$f(x, y) = c(x^2 - y^2)e^{-x}$$

for $0 \leq x < \infty, -x \leq y \leq x$. For $x > 0$, we have

$$f_X(x) = \int_{-x}^x c(x^2 - y^2)e^{-x} dy = \frac{4c}{3}x^3e^{-x}.$$

Hence, $f_{Y|X}(y|x) = \frac{3}{4} \frac{x^2 - y^2}{x^3}$ for $-x < y < x$, 0 otherwise. We conclude that

$$F_{Y|X}(y|x) = \begin{cases} 0 & y \leq -x \\ \frac{3}{4} \int_{-x}^y \frac{x^2 - y^2}{x^3} dy = \frac{1}{4} \left(\frac{y(3x^2 - y^2)}{x^3} + 2 \right) & -x < y < x \\ 1 & x \leq y. \end{cases}$$